

WAY TO SUCCESS PUBLICATIONS
HALF - YEARLY EXAMINATION
ANSWER KEY
10TH MATHEMATICS

Part - I

Answer all the questions	
1.	c) {4, 9, 25, 49, 121}
2.	c) $\frac{2}{9x^2}$
3.	b) 2
4.	b) An arithmetic progression
5.	c) $\frac{x^2-7x+40}{(x^2-25)(x+1)}$
6.	b) $\begin{bmatrix} 2 & 2 \\ 2 & -1 \end{bmatrix}$
7.	c) 4
8.	b) 25 sq.units
9.	b) $x + y = 3, 3x + y = 7$
10.	d) 1
11.	b) 4 cm
12.	c) 3π
13.	b) 100
14.	b) $\frac{7}{10}$

Part - II

15. (i) Range of $f = \{1, 8, 27, 64\}$
(ii) Since distinct elements in A are mapped into distinct images in B , it is a one-one function. $2 \in B$ is not the image of any element of A . So, it is Into function.
16. $800 = 2 \times 2 \times 2 \times 2 \times 2 \times 5 \times 5$
 $= 2^5 \times 5^2$
 $a^b \times b^a = 2^5 \times 5^2$
 $a = 5$ and $b = 2$.
17. $a_n = \frac{1}{3}n + \frac{1}{6}$
 $a_n = \frac{2}{6}n + \frac{3}{6} - \frac{2}{6}$
 $a_n = \frac{3}{6} + (n-1)\frac{2}{6}$
It is in the form $a_n = a + (n-1)d$.
Here $a = \frac{3}{6}$, $d = \frac{2}{6}$
Given sequence is an A.P

$$18. n = 27 + 1 = 28$$

$$1 + 3 + 5 + \dots + 55 = (28)^2 = 784.$$

$$19. \alpha + \beta = -6, \alpha\beta = -4$$

$$(\alpha - \beta)^2 = (\alpha + \beta)^2 - 4\alpha\beta$$

$$= (-6)^2 - 4(-4)$$

$$= 36 + 16$$

$$= 52$$

$$20. A^T = \begin{bmatrix} 5 & -\sqrt{7} & 8 \\ 2 & 0.7 & 3 \\ 2 & \frac{5}{2} & 1 \end{bmatrix}$$

$$(A^T)^T = \begin{bmatrix} 5 & 2 & 2 \\ -\sqrt{7} & 0.7 & \frac{5}{2} \\ 8 & 3 & 1 \end{bmatrix} = A$$

$$21. \text{Slope } m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{3-a}{9-(-2)} = \frac{3-a}{11}$$

$$\frac{3-a}{11} = -\frac{1}{2} \text{ (Given)}$$

$$a = \frac{17}{2}$$

22. Area of the triangle

Area of the triangle

$$= \frac{1}{2} \begin{vmatrix} 1 & -4 & -3 & 1 \\ -1 & 6 & -5 & -1 \end{vmatrix}$$

$$= \frac{1}{2} [29 + 19] = \frac{1}{2} (48) = 24 \text{ Sq. units}$$


$$23. \tan \theta = \frac{AB}{BC} = \frac{10\sqrt{3}}{30}$$

$$\tan \theta = \frac{1}{\sqrt{3}}$$

$$\tan 30^\circ = \frac{1}{\sqrt{3}}$$

$$\therefore \theta = 30^\circ$$

$\therefore$ The angle of elevation $\theta = 30^\circ$



$$24. \text{Given that, } \frac{r_1}{r_2} = \frac{12}{16} = \frac{3}{4}$$

$$\text{Ratio of C.S.A. of balloons} = \frac{4\pi r_1^2}{4\pi r_2^2} = \frac{9}{16}$$

Ratio of C.S.A. of balloons is 9:16.

$$25. \text{Volume of the cone} = 11088 \text{ cm}^3$$

$$\frac{1}{3} \pi r^2 h = 11088$$

$$\frac{1}{3} \times \frac{22}{7} \times r^2 \times 24 = 11088$$

$$r^2 = 441$$

Therefore, radius of the cone $r = 21 \text{ cm}$.

$$26. \pi r^2 h = \frac{4}{3} \pi r^3$$

$$\pi \times 10 \times 10 \times h = \frac{4}{3} \times \pi \times 15 \times 15 \times 15$$

$$h = 45 \text{ cm}$$

Height of the cylinder = 45 cm

$$27. R = 36.8, S = 13.4, L = ?$$

$$R + S = L$$

$$L = 36.8 + 13.4$$

$$L = 50.2$$

$$28. S = \{HH, HT, TH, TT\}$$

$$n(S) = 4$$

$$P(A) = \frac{n(A)}{n(S)} = \frac{2}{4} = \frac{1}{2}$$

Part - III

$$29. A = \{2,3\}, B = \{0,1\}, C = \{1,2\}$$

$$B \cap C = \{1\}$$

$$A \times (B \cap C) = \{2,3\} \times \{1\}$$

$$A \times (B \cap C) = \{(2,1), (3,1)\} \dots (1)$$

$$A \times C = \{(2,1), (2,2), (3,1), (3,2)\}$$

$$(A \times B) \cap (A \times C) = \{(2,1), (3,1)\} \dots (2)$$

From (1) and (2),

$$A \times (B \cap C) = (A \times B) \cap (A \times C)$$

$$30. f \circ g(x) = f(6x - k) = 18x - 3k + 2$$

$$g \circ f(x) = g(3x + 2) = 18x + 12 - k$$

Given that, $f \circ g = g \circ f$

$$18x - 3k + 2 = 18x + 12 - k$$

$$k = -5$$

$$31. t_4 = 8 \Rightarrow ar^3 = 8 \dots (1)$$

$$t_8 = \frac{128}{625} \Rightarrow ar^7 = \frac{128}{625} \dots (2)$$

$$(2) \div (1) \Rightarrow r^4 = \left(\frac{2}{5}\right)^4 \Rightarrow r = \frac{2}{5}$$

$$\text{From (1)} \Rightarrow a \left(\frac{2}{5}\right)^3 = 8 \Rightarrow a = 125$$

Required G.P, $a, ar, ar^2, \dots$

125, 50, 20, ...

$$32. 10^2 + 11^2 + 12^2 + \dots + 24^2$$

$$= (1^2 + 2^2 + \dots + 24^2)$$

$$-(1^2 + 2^2 + 3^2 + \dots + 9^2)$$

$$= \frac{24(24+1)(24 \times 2 + 1)}{6} - \frac{9(9+1)(2 \times 9 + 1)}{6}$$

$$= 4(25)(49) - 3(5)(19)$$

$$= 4900 - 285$$

$$= 4615 \text{ cm}^2$$

$$33. x + y + z = 5 \dots (1)$$

$$2x - y + z = 9 \dots (2)$$

$$x - 2y + 3z = 16 \dots (3)$$

Add (1) + (2)

$$(1) \Rightarrow x + y + z = 5$$

$$(2) \Rightarrow 2x - y + z = 9$$

$$\frac{3x}{+ 2z} = 14 \dots (4)$$

$$(2) \times 2 \Rightarrow 4x - 2y + 2z = 18$$

$$(3) \Rightarrow x - 2y + 3z = 16$$

$$\begin{array}{r} (-) \quad (+) \quad (-) \quad (-) \\ 3x \quad \quad - z = 2 \dots (5) \end{array}$$

$$(4) \Rightarrow 3x + 2z = 14$$

$$(5) \Rightarrow 3x - z = 2$$

$$\begin{array}{r} (-) \quad (+) \quad (-) \\ 3z = 12 \end{array}$$

$$3z = 12$$

$$z = \frac{12}{3} = 4$$

$z = 4$ Substituting in (5)

$$3x - 4 = 2 \Rightarrow 3x = 6 \Rightarrow x = \frac{6}{3} \Rightarrow x = 2$$

$x = 2, z = 4$ Substituting in (1)

$$y = 5 - 2 - 4 \Rightarrow y = -1$$

$$\therefore x = 2, y = -1, z = 4$$

34.

$$\begin{array}{r} 3x^2 + 2x + 4 \\ 3x^2 \overline{) 9x^2 + 12x^3 + 28x^2 + ax + b} \\ \underline{9x^2} \\ 6x^2 + 2x \\ \underline{6x^2 + 2x} \\ 6x^2 + 4x + 4 \\ \underline{6x^2 + 4x + 4} \\ 0 \end{array}$$

Given polynomial is a perfect square

$$a - 16 = 0, b - 16 = 0$$

Therefore $a = 16, b = 16$

35. $n(S) = 36$

$$A = \{(1,1), (2,2), (3,3), (4,4), (5,5), (6,6)\}$$

$$B = \{(1,3), (2,2), (3,1)\}$$

$$A \cap B = \{(2,2)\}$$

$$n(A) = 6, n(B) = 3, n(A \cap B) = 1$$

$$P(A) = \frac{6}{36}, P(B) = \frac{3}{36}$$

$$P(A \cap B) = \frac{n(A \cap B)}{n(S)} = \frac{1}{36}$$

$$P(A \cup B) = \frac{6}{36} + \frac{3}{36} - \frac{1}{36} = \frac{8}{36} = \frac{2}{9}$$

36. $A - B = \begin{bmatrix} -3 & 2 \\ 0 & -2 \end{bmatrix}$

$$(A - B)^T = \begin{bmatrix} -3 & 2 \\ 0 & -2 \end{bmatrix} \dots \dots \dots (1)$$

$$A^T - B^T = \begin{bmatrix} 1 & 1 \\ 2 & 3 \end{bmatrix} - \begin{bmatrix} 4 & 1 \\ 0 & 5 \end{bmatrix} = \begin{bmatrix} -3 & 0 \\ 2 & -2 \end{bmatrix} \dots (2)$$

$$\text{From (1) and (2), } (A - B)^T = A^T - B^T$$

37.

Pythagoras Theorem


Statement: In a right angle triangle, the square on the hypotenuse is equal to the sum of the squares on the other two sides.

Proof:

Given : In $\triangle ABC$, $\angle A = 90^\circ$

To prove : $AB^2 + AC^2 = BC^2$

Construction : Draw $AD \perp BC$



No.	Statement	Reason
1.	Compare $\triangle ABC$ and $\triangle ABD$ $\angle B$ is common $\angle BAC = \angle BDA = 90^\circ$ Therefore, $\triangle ABC \sim \triangle ABD$ $\frac{AB}{AD} = \frac{BC}{AB}$ $AB^2 = BC \times BD$ $\dots \dots (1)$	Given $\angle BAC = 90^\circ$ and by construction $\angle BDA = 90^\circ$ By AA similarity
2.	Compare $\triangle ABC$ and $\triangle ADC$ $\angle C$ is common $\angle BAC = \angle ADC = 90^\circ$ Therefore, $\triangle ABC \sim \triangle ADC$ $\frac{BC}{AC} = \frac{AC}{DC}$ $AC^2 = BC \times DC$ $\dots (2)$	Given $\angle BAC = 90^\circ$ and by construction $\angle CDA = 90^\circ$ By AA similarity

Adding (1) and (2) we get


$$AB^2 + AC^2 = BC \times BD + BC \times DC$$

$$= BC (BD + DC) = BC \times BC$$

$$AB^2 + AC^2 = BC^2$$

Hence the theorem is proved.

38.



$$\text{Slope of } BC = \frac{y_2 - y_1}{x_2 - x_1} = \frac{3 - (-2)}{12 - 10} = \frac{5}{2}$$

Equation of the altitude passing through the vertex A.

$$y - y_1 = m(x - x_1)$$

$$A(-3, 0) \text{ and } m = \frac{5}{2}$$

$$y - 0 = -\frac{1}{\frac{5}{2}}(x - (-3))$$

$$y = -\frac{2}{5}(x + 3)$$

$$5y = -2x - 6$$

$$2x + 5y + 6 = 0$$

39. $\tan 45^\circ = \frac{30}{AB}$

$$AB = 30m$$

$$\tan 60^\circ = \frac{30+h}{30}$$

$$30\sqrt{3} = 30 + h$$


$$h = 30(\sqrt{3} - 1)$$

$$= 30(1.732 - 1)$$

$$= 30(0.732)$$

$$= 21.96$$

The height of the tower is 21.96 m.


 40. Hemisphere : $r = 7$

Cone: $r = 7, l = 11$

$$\text{Surface area of doll} = 2\pi r^2 + \pi r l$$

$$= \left(2 \times \frac{22}{7} \times 7 \times 7\right) + \left(\frac{22}{7} \times 7 \times 11\right)$$

$$= \frac{22}{7} \times 7(14 + 11)$$

$$= 550cm^2$$

41.

x	f	$d = x - A$	fd	fd^2
10	3	-8	-24	192
15	2	-3	-6	18
18	5	0	0	0
20	8	2	16	32
25	2	7	14	98
$N = 20$			0	340

$$\sigma = \sqrt{\frac{\sum f_i d_i^2}{N} - \left(\frac{\sum f_i d_i}{N}\right)^2} = \sqrt{\frac{340}{20}} = \sqrt{17} \cong 4.1$$

42. $\Delta = b^2 - 4ac$

$$(b - c)^2 - 4(a - b)(c - a) = 0$$


$$b^2 + c^2 + 4a^2 + 2bc - 4ac - 4ab = 0$$

$$(2a - b - c)^2 = 0$$

$$2a = b + c$$


Part - IV

43. a)



b)

Rough diagram




Verification: $PT = \sqrt{OP^2 - OT^2}$

$$= \sqrt{(7.2)^2 - (3.6)^2} = \sqrt{38.7} \cong 6.2 \text{ cm.}$$

44. a) $x^2 - 9x + 20 = 0$

$$y = x^2 - 9x + 20$$

x	-3	-2	-1	0	1	2	3	4	5	6
x^2	9	4	1	0	1	4	9	16	25	36
$-9x$	27	18	9	0	-9	-18	-27	-36	-45	-54
20	20	20	20	20	20	20	20	20	20	20
y	56	42	30	20	12	6	2	0	0	2

 Points: $(-3, 56), (-2, 42), (-1, 30), (0, 20), (1, 12), (2, 6), (3, 2), (4, 0), (5, 0), (6, 2)$


Solution $x = \{4, 5\}$

Real and unequal roots.

44.b) $y = x^2 + x - 2$

x	-3	-2	-1	0	1	2
y	4	0	-2	-2	0	4

Solve

$$y = x^2 + x - 2$$

$$0 = x^2 + x - 2 \quad (-)$$

$$y = 0$$

